arXiv:math-ph/9908001v1 1 Aug 1999

A new differential calculus on noncommutative spaces

R. P. Malik [1]
S. N. Bose National Centre for Basic Sciences,
Block-JD, Sector-III, Salt Lake, Calcutta-700 091, India.

A. K. Mishra [2] and **G. Rajasekaran** [3]
The Institute of Mathematical Sciences,
C.I.T Campus, Madras-600 113, India.

Abstract

We develop a $GL_{qp}(2)$ invariant differential calculus on a two-dimensional noncommutative quantum space. Here the co-ordinate space for the exterior quantum plane is spanned by the differentials that are commutative (bosonic) in nature.

The quantum group $GL_{qp}(n)$ covariant differential calculus on a quantum hyperplane is generated by three elements : (i) $\{x_i\}(i = 1, 2, 3, ...n)$, the noncommuting co-ordinates of the quantum plane, (ii) differentials $\{dx_i\}$, which are co-ordinates of the exterior quantum plane, and (iii) derivatives on the quantum plane, namely $\{\frac{\partial}{\partial x_i}\}$ [1-4]. In addition, one also introduces an exterior differential d [1]

$$d \;=\; \sum_i dx_i \frac{\partial}{\partial x_i}. \tag{1}$$

In the classical limit, two distinct kinds of dx, satisfying either the algebraic relation

$$\tilde{d}x_i \tilde{d}x_j + \tilde{d}x_j \tilde{d}x_i \;=\; 0 \tag{2}$$

or the relation

$$dx_i dx_j \;-\; dx_j dx_i \;=\; 0 \tag{3}$$

exist. The one-form $\tilde{d}x$, with $(\tilde{d}x)^2 = 0$, is introduced in the context of differential geometry [5]. On the other hand, dx denotes the infinitesimal whose square is not identically zero. The co-ordinate space for the exterior plane, spanned by $\{\tilde{d}x_i\}$, is anticommutative (fermionic), and the one spanned by $\{dx_i\}$ is commutative (bosonic).

---

[1] E-mail address: malik@boson.bose.res.in
[2] E-mail address: mishra@imsc.ernet.in
[3] E-mail address: graj@imsc.ernet.in

Much progress has been made in the area of noncommutative differential calculus (NDC) based on a generalized $\tilde{d}x$ [1-4,6,7]. Is it possible to provide a consistent noncommutative differential calculus involving appropriately modified dx? The answer to this query is in affirmative. In the present paper, we develop the basic structure of this new calculus with a view to evaluate the derivatives of a function having as its arguments the co-ordinates of quantum plane and/or exterior quantum plane (i.e., differentials), or any product of such functions.

We denote henceforth the conventional noncommutative differential calculus having $\tilde{d}x_i$ as one of its elements as NDC-I, and the new differential calculus involving dx_i as NDC-II. The d and dx_i in eq.(1) now belong to NDC-II, whereas these quantities are to be replaced by $\tilde{d}$ and $\tilde{d}x_i$ in NDC-I. It can be shown that $\tilde{d}^2 = 0$ [1], though no such constraint exists for d^2. In fact, $d^n \neq 0$ for any integer $n = 0, 1, 2, \ldots$. Consequently, we have non-zero $d^n x_i$ - the nth differential in NDC-II, with $d^0 x_i \equiv x_i$. In the NDC-I, all higher order differentials, but for the one corresponding to $n = 1$, are zero. Here again, $\tilde{d}^0 x_i \equiv x_i$.

The derivatives $\{\frac{\partial}{\partial(\tilde{d}x_i)}\}$ on the exterior quantum plane, though not defined originally, has been subsequently introduced [8]. We presently introduce the analogous derivatives $\{\frac{\partial}{\partial(d^n x_i)}\}$. How these additional derivatives modify d in eq.(1) will be considered shortly.

We consider here a two dimensional system. This enables us to describe the basic structure of the new calculus without getting involved in the mathematical complexities when dimension exceeds two. The case of higher dimensions will be considered in a subsequent paper. We identify $d^n x_1 \equiv d^n x$ and $d^n x_2 \equiv d^n y$. In order to provide the details of NDC-II, the first task is to determine the generalized commutation relations between $(d^n x, d^m x), (d^n y, d^m y)$ and $(d^n x, d^m y)$ for arbitrary positive integers n and m. In addition, either or both n and m can be zero also. To achieve this, one needs only the relations corresponding to $m = n, n+1$. The remaining relations follow by the successive application of the exterior differential d. The d satisfies the Leibnitz rule

$$d\,(fg) = (df)\,g + f\,(dg) \tag{4}$$

with f and g being functions of $(d^n x, d^m y)$.

We begin with the following general relations

$$(d^n x)(d^n y) = C_1 (d^n y)(d^n x) \tag{5}$$

$$(d^n x)(d^{n+1} x) = C_2 (d^{n+1} x)(d^n x) \tag{6}$$

$$(d^n y)(d^{n+1} y) = C_3 (d^{n+1} y)(d^n y) \tag{7}$$

$$(d^n y)(d^{n+1} x) = C_4 (d^{n+1} x)(d^n y). \tag{8}$$

The various C'^s are non-zero constants. Applying d on relation (5) and using eq.(8), we obtain the last relation

$$(d^n x)(d^{n+1} y) \;=\; C_1 (d^{n+1} y)(d^n x) + (C_1 C_4 - 1)(d^{n+1} x)(d^n y). \tag{9}$$

The requirement of associativity and the validity of Yang-Baxter equation demands that when $d^k x, d^k y$ $(k = n, n+1)$ are commuted through eqs. (5-8), either from the left or from the right, no new relation should emerge [9]. This condition is satisfied provided

$$C_2 \;=\; C_3 \;= C_1 C_4. \tag{10}$$

It can be seen that when $C_1 \;=\; q$ and $C_4 \;=\; p$, the relations (5-8) are invariant under the quantum group transformations

$$\begin{pmatrix} d^n x \\ d^n y \end{pmatrix} \rightarrow \begin{pmatrix} d^n x' \\ d^n y' \end{pmatrix} = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \begin{pmatrix} d^n x \\ d^n y \end{pmatrix} \tag{11}$$

where A, B, C, D are the elements of a 2×2 $GL_{qp}(2)$ matrix obeying the q-algebraic relations in rows and columns as

$$\begin{array}{lclcl} AB = pBA & , & AC = qCA & , & BC = \frac{q}{p} CB \\ CD = pDC & , & BD = qDB & , & AD - DA = (p - q^{-1}) BC \end{array} \tag{12}$$

and these elements are assumed to commute with all the differentials in eqs.(5-8).

Substituing the values of C's in eqs. (5-9), we get

$$\xi^n \eta^n \;=\; q \eta^n \xi^n \tag{13}$$

$$\xi^n \xi^{n+1} \;=\; pq \xi^{n+1} \xi^n \tag{14}$$

$$\eta^n \eta^{n+1} \;=\; pq \eta^{n+1} \eta^n \tag{15}$$

$$\eta^n \xi^{n+1} = p \xi^{n+1} \eta^n \tag{16}$$

$$\xi^n \eta^{n+1} = q \eta^{n+1} \xi^n + (pq-1) \xi^{n+1} \eta^n \tag{17}$$

where for the notational convenience, we have introduced

$$d^k x \equiv \xi^k \quad ; \quad d^k y \equiv \eta^k. \tag{18}$$

We note here that superscripts of ξ and η denote an index, and not the exponent.

Applying the d consecutively on relations (13-17), and after some algebraic manipulations, the following generalizations are obtained for equations (14-17) corresponding to any $m > n$:

$$\xi^n \xi^m = pq \xi^m \xi^n + (pq-1) \sum_{\substack{r=1 \\ m-n-r>0}}^{m-n-1} \xi^{n+r} \xi^{m-r} \tag{19}$$

$$\eta^n \eta^m = pq \eta^m \eta^n + (pq-1) \sum_{\substack{r=1 \\ m-n-r>0}}^{m-n-1} \eta^{n+r} \eta^{m-r} \tag{20}$$

$$\eta^n \xi^m = p \xi^m \eta^n + (pq-1) \sum_{\substack{r=1 \\ m-n-r>0}}^{m-n-1} \eta^{m-r} \xi^{n+r} \tag{21}$$

$$\xi^n \eta^m = q \eta^m \xi^n + (pq-1) \sum_{\substack{r=0 \\ m-n-r>0}}^{m-n-1} \xi^{m-r} \eta^{n+r}. \tag{22}$$

Equations (13,19-22) provide all possible commutation relations between the differentials.

The ultimate aim of the present analysis is to provide relations which enable one to evaluate the derivatives of functions like f and g introduced in eq.4. Such derivatives can be obtained using the commutation relations between (ξ^n, η^m) and derivatives $(\frac{\partial}{\partial \xi^k}, \frac{\partial}{\partial \eta^l})$. To obtain these relations, we need explicit form of the exterior differential d. Accordingly, we generalize the expression (1) as

$$d = \sum_{n=1}^{\infty} \left(\xi^n \frac{\partial}{\partial \xi^{n-1}} + \eta^n \frac{\partial}{\partial \eta^{n-1}} \right). \tag{23}$$

Applying d on $\xi^n f$ and using (23), we obtain

$$d(\xi^n f) = \sum_{m>0} \xi^m \frac{\partial}{\partial \xi^{m-1}} \xi^n f + \sum_{m>0} \eta^m \frac{\partial}{\partial \eta^{m-1}} \xi^n f. \tag{24}$$

The Leibnitz rule (cf. eq.4) leads to

$$d\,(\xi^n f) = \xi^{n+1}\, f + \xi^n\, d\, f \tag{25}$$

where $\xi^{n+1} = d\xi^n = d(d^n x) = d^{n+1}x$. Substituting d from eq.(23) in the right hand side of eq.(25), we get

$$d(\xi^n f) \;=\; \xi^{n+1} f + \xi^n \sum_{m>0} \xi^m \frac{\partial}{\partial \xi^{m-1}} f + \xi^n \sum_{m>0} \eta^m \frac{\partial}{\partial \eta^{m-1}} f. \tag{26}$$

Next, we use relations (13,19-22) to exchange the order of $\xi^n \xi^m$ and $\xi^n \eta^m$ in eq.(26). Thereafter, comparing the coefficients of ξ^m and η^m in the resulting expression with the coefficients of ξ^m and η^m in eq.(24), we obtain the following set of required commutation relations

$$\frac{\partial}{\partial \xi^m}\xi^n \;=\; \frac{1}{pq}\xi^n \frac{\partial}{\partial \xi^m} - \frac{(pq-1)}{pq}\sum_{k=1}^{m} \xi^{n-k}\frac{\partial}{\partial \xi^{m-k}}, \quad 0 \le m \le n-2 \tag{27}$$

$$\frac{\partial}{\partial \xi^m}\xi^n \;=\; \xi^n \frac{\partial}{\partial \xi^m}, \qquad m = n-1 \tag{28}$$

$$\frac{\partial}{\partial \xi^m}\xi^n \;= \delta_{n,m} + \; pq\xi^n \frac{\partial}{\partial \xi^m} + (pq-1)\sum_{k=1}^{\infty}\left(\xi^{n+k}\frac{\partial}{\partial \xi^{m+k}} + \eta^{n+k-1}\frac{\partial}{\partial \eta^{m+k-1}}\right), \; m > n-1 \tag{29}$$

$$\frac{\partial}{\partial \eta^m}\xi^n \;=\; \frac{1}{p}\xi^n \frac{\partial}{\partial \eta^m} - \frac{(pq-1)}{p}\sum_{k=1}^{m} \xi^{n-k}\frac{\partial}{\partial \eta^{m-k}}, \quad 0 \le m \le n-2 \tag{30}$$

$$\frac{\partial}{\partial \eta^m}\xi^n \;=\; q\xi^n \frac{\partial}{\partial \eta^m}, \quad m \ge n-1. \tag{31}$$

Repeating the same procedure with $d(\eta f)$ in the place of $d(\xi f)$, we get

$$\frac{\partial}{\partial \eta^m}\eta^n \;=\; \frac{1}{pq}\eta^n \frac{\partial}{\partial \eta^m} - \frac{(pq-1)}{pq}\sum_{k=1}^{m} \eta^{n-k}\frac{\partial}{\partial \eta^{m-k}}, \quad 0 \le m \le n-2 \tag{32}$$

$$\frac{\partial}{\partial \eta^m}\eta^n \;=\; \eta^n \frac{\partial}{\partial \eta^m}, \quad m = n-1 \tag{33}$$

$$\frac{\partial}{\partial \eta^m}\eta^n \;= \delta_{n,m} + \; pq\eta^n \frac{\partial}{\partial \eta^m} + (pq-1)\sum_{k=1}^{\infty}\left(\xi^{n+k}\frac{\partial}{\partial \xi^{m+k}} + \eta^{m+k}\frac{\partial}{\partial \eta^{m+k}}\right), \quad m > n-1 \tag{34}$$

$$\frac{\partial}{\partial \xi^m}\eta^n \;=\; \frac{1}{q}\eta^n \frac{\partial}{\partial \xi^m} - \frac{(pq-1)}{q}\sum_{k=1}^{m} \eta^{n-k}\frac{\partial}{\partial \xi^{m-k}}, \quad 0 \le m \le n-2 \tag{35}$$

$$\frac{\partial}{\partial \xi^m}\eta^n \;=\; \frac{1}{q}\eta^n \frac{\partial}{\partial \xi^m} - \frac{(pq-1)}{q}\sum_{k=1}^{n-1} \eta^{k}\frac{\partial}{\partial \xi^{k-1}}, \quad m = n-1 \tag{36}$$

$$\frac{\partial}{\partial \xi^m}\eta^n \;=\; p\eta^n \frac{\partial}{\partial \xi^m}, \quad m > n-1. \tag{37}$$

Given any function of (ξ^n, η^m), the relations (27-37) enable one to evaluate its derivative as well as any sequence of derivatives with respect to ξ^k and η^l.

We may note the importance of the new calculus in the context of particle dynamics on a noncommutative space. For example, it is the dx and d^2x which are related to the velocity $\dot{x}$ and acceleration $\ddot{x}$ through the relations $dx = \dot{x}dt$ and $d^2x = \ddot{x}(dt)^2$, where the time t is taken to be a commuting number. On the other hand, a relation like $\tilde{d}x = \dot{x}dt$ would make the velocity nilpotent, and the acceleration can not be related to $\tilde{d}^2x$ as this quantity is identically zero.

Is it possible to provide a connection between NDC-I and the new noncommutative calculus? Can the new calculus be mapped to a Fock-space structure? These queries along with the possible applications will be considered in our future communications.

References

1. Wess J and Zumino B 1990 *Nucl. Phys. (Proc. Suppl.)* **B18** 302
2. Manin Yu I 1989 *Comm. Math. Phys.* **123** 163
3. Woronowicz S L 1989 *Comm. Math. Phys.* **122** 125
4. Schirrmacher A 1991 *Z. Phys.* **C50** 321
5. Spivak M 1970 *A comprehensive Introduction to Differential Geometry* (Publish or Perish, Boston) Vol. **1**
6. Pusz W and Woronowicz S L 1989 *Rep. Math. Phys.* **27** 231
7. Soni S K 1991 *J. Phys.* **A24** L169
8. Mishra A K and Rajasekaran G. 1997 *J. Math. Phys.* **38** 466
9. Lukin M, Stern A and Yakushin I 1993 *J. Phys.* **A26** 5115